

Serre's conjecture over $\mathbb{F}_9$

Jordan S. Ellenberg

10 May 2001

Abstract

In this paper we show that an odd Galois representation $\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{F}_9)$ satisfying certain local conditions at 3 and 5 is modular. Our main tool is an idea of Taylor [12], which reduces the problem to that of exhibiting points on a Hilbert modular surface which are defined over a solvable extension of $\mathbb{Q}$, and which satisfy certain reduction properties. As a corollary, we show that Hilbert-Blumenthal abelian surfaces with good ordinary reduction at 3 and 5 are modular.

Introduction

In 1986, J. P. Serre proposed the following conjecture:

Conjecture. *Let $\mathbb{F}$ be a finite field of characteristic p , and*

$$\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{F})$$

an irreducible representation such that $\det \bar{\rho}$ applied to complex conjugation yields -1 . Then $\bar{\rho}$ is the mod p representation attached to a modular form on $\text{GL}_2(\mathbb{Q})$.

Serre's conjecture, if true, would provide the first serious glimpse into the non-abelian structure of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. The work of Langlands and Tunnell shows that Serre's conjecture is true when $\text{GL}_2(\mathbb{F})$ is solvable; that is, when $\mathbb{F}$ is $\mathbb{F}_2$ or $\mathbb{F}_3$. Work of Shepherd-Barron and Taylor [9] and Taylor [12] have shown that the conjecture is also true, under some local conditions on $\bar{\rho}$, when $\mathbb{F}$ is $\mathbb{F}_4$ or $\mathbb{F}_5$.

In the present work, we show that Serre's conjecture is true, again subject to certain local conditions, when $\mathbb{F} = \mathbb{F}_9$. To be precise, we prove the following theorem.

Theorem. *Let*

$$\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{F}_9)$$

be an odd Galois representation such that

- *The restriction of $\bar{\rho}$ to D_3 can be written as*

$$\bar{\rho}|_{D_3} \cong \begin{bmatrix} \psi_1 & * \\ 0 & \psi_2 \end{bmatrix},$$

where ψ_1 and ψ_2 are characters such that $(\psi_1 \oplus \psi_2)|_{I_3}$ is isomorphic to the direct sum of the trivial character and $\bar{\chi}_3$;

- The image of the inertia group I_5 lies in $\mathrm{SL}_2(\mathbb{F}_9)$, and has odd order in $\mathrm{PSL}_2(\mathbb{F}_9)$.

Then $\bar{\rho}$ is modular.

As a corollary, we get the following result towards a generalized Shimura-Taniyama-Weil conjecture for abelian surfaces:

Corollary. *Let $A/\mathbb{Q}$ be a Hilbert-Blumenthal abelian surface which has good ordinary reduction at 3 and 5. Then A is a quotient of $J_0(N)$ for some integer N .*

We prove the theorem above by exhibiting $\bar{\rho}$ as the Galois representation on the 3-torsion subscheme of a certain Hilbert-Blumenthal abelian surface defined over a totally real extension $F/\mathbb{Q}$ with solvable Galois group. We then use an idea of Taylor, together with a new theorem of Skinner and Wiles [11], to prove the modularity of the abelian surface, and consequently of $\bar{\rho}$.

The key algebro-geometric point is that a certain twisted Hilbert modular variety has many points defined over solvable extensions of $\mathbb{Q}$. This suggests that we consider the class of varieties X such that, if K is a number field, and Σ is the set of all solvable Galois extensions L/K , then

$$\bigcup_{L \in \Sigma} X(L)$$

is Zariski-dense in X . We say X has “property **S**” in this case. Certainly if X has a Zariski-dense set of points over a *single* number field—for example, if X is unirational—it has property **S**. The Hilbert modular surfaces we consider, on the other hand, are varieties of general type with property **S**.

To indicate our lack of knowledge about solvable points on varieties, note that at present there does not exist a variety which we can prove does *not* have property **S**! Nonetheless, it seems reasonable to guess that “sufficiently complicated” varieties do not have property **S**.

One might consider the present result evidence for the truth of Serre’s conjecture. On the other hand, it should be pointed out that the theorems here and in [9], [12] rely crucially on the facts that

- the GL_2 of small finite fields is solvable, and
- certain Hilbert modular varieties for number fields of small discriminant have property **S**.

These happy circumstances may not persist very far. In particular, it is reasonable to guess that only finitely many Hilbert modular varieties have property **S**. If so, one might say that we have much philosophical but little numerical evidence for the truth of Serre’s conjecture in general. Our ability to compute has progressed mightily since Serre’s conjecture was first announced. It would be interesting, given the present status of the conjecture, to carry out numerical experiments for $\mathbb{F}$ a “reasonably large” finite field—whatever that might mean.

Notation

If $\bar{\rho} : \mathrm{Gal}(\bar{K}/K) \rightarrow \mathrm{GL}_2(\mathbb{F}_9)$ is a Galois representation, we define $V_{\bar{\rho}}$ to be the symplectic Galois module $\mathbb{F}_9 \oplus \mathbb{F}_9$, with Galois acting by $\bar{\rho}$, and the standard symplectic pairing.

If K is a totally real number field, write $c \in \mathrm{Gal}(\bar{K}/K)$ for a complex conjugation. If v is a prime of a number field F , we write $D_v \subset G_F$ for the decomposition group associated to v , and I_v for the corresponding inertia group. The p -adic cyclotomic character of Galois is denoted by χ_p , and its mod p reduction by $\bar{\chi}_p$.

If $V \subset \mathbb{P}^N$ is a projective variety, write $F_1(V)$ for the Fano variety of lines contained in V .
If $\mathcal{O}$ is a ring, an $\mathcal{O}$ -module scheme is an $\mathcal{O}$ -module in the category of schemes.
All Hilbert modular forms are understood to be of parallel weight.

1 Realizations of Galois representations on HBAV's

Recall that a *Hilbert-Blumenthal abelian variety* (HBAV) over a number field is an abelian d -fold endowed with an injection $\mathcal{O} \hookrightarrow \text{End}(A)$, where $\mathcal{O}$ is the ring of integers of a totally real number field of degree d over $\mathbb{Q}$. Many Hilbert-Blumenthal abelian varieties can be shown to be modular; for example, see [9]. It is therefore sometimes possible to show that a certain mod p Galois representation $\bar{\rho}$ is modular by realizing it on the p -torsion subscheme of some HBAV.

We will show that, given a Galois representation $\text{Gal}(\bar{K}/K) \rightarrow \text{GL}_2(\mathbb{F}_9)$ satisfying some local conditions at 3, 5 and ∞ , we can find an abelian surface over a solvable extension of K , satisfying some local conditions at 3 and 5. One of these conditions—that certain representations be “ D_p -distinguished”—requires further comment.

Definition 1.1. Let $\bar{\rho} : \text{Gal}(\bar{K}/K) \rightarrow \text{GL}_2(\bar{\mathbb{F}}_p)$ be a Galois representation, and let $\mathfrak{p}|p$ be a prime of K . We say that $\bar{\rho}$ is $D_{\mathfrak{p}}$ -distinguished if the semisimplification of the restriction $\text{Gal}(\bar{K}/K)|_{D_{\mathfrak{p}}}$ is isomorphic to $\theta_1 \oplus \theta_2$, with θ_1 and θ_2 distinct characters from $\text{Gal}(\bar{K}/K)$ to $\bar{\mathbb{F}}_p^*$.

This condition is useful in deformation theory, and is required, in particular, in the main theorem of [11]. A natural source of $D_{\mathfrak{p}}$ -distinguished Galois representations is provided by abelian varieties with ordinary reduction at $\mathfrak{p}$.

Proposition 1.2. *Let p be an odd prime. Let K_v be a finite extension of $\mathbb{Q}_p$ with odd ramification degree, and let A/K_v be a HBAV with good ordinary or multiplicative reduction and real multiplication by $\mathcal{O}$, and let $\mathfrak{p}$ be a prime of $\mathcal{O}$ dividing p .*

Then the semisimplification of the $\text{Gal}(\bar{K}_v/K_v)$ -module $A[\mathfrak{p}]$ is isomorphic to $\theta_1 \oplus \theta_2$, with θ_1 and θ_2 distinct characters of $\text{Gal}(\bar{K}_v/K_v)$.

Proof. We think of $A[\mathfrak{p}]$ as a two-dimensional $\mathcal{O}/\mathfrak{p}\mathcal{O}$ -module, where $\mathcal{O}$ is the ring of real multiplications. Let e be the ramification index of K_v over $\mathbb{Q}_p$.

The decomposability of $A[\mathfrak{p}]$ into an extension of θ_1 by θ_2 is a general fact about ordinary abelian varieties.

Now the action of the inertia group I_v on $A[\mathfrak{p}]$ is an extension of the trivial character by the cyclotomic character. Since the ramification index of $K_v/\mathbb{Q}_p$ is odd, these two characters are distinct. Therefore, θ_1 and θ_2 are also distinct. $\square$

We are now ready to state the main theorem of this section.

Proposition 1.3. *Let K be a totally real number field, and let*

$$\bar{\rho} : \text{Gal}(\bar{K}/K) \rightarrow \text{GL}_2(\mathbb{F}_9)$$

be a Galois representation such that $\det \bar{\rho} = \bar{\chi}_3$. Suppose that

- *The absolute ramification degree of K is odd at every prime of K above 3 and 5.*

- For any prime w of K over 3, the restriction of $\bar{\rho}$ to the decomposition group D_w can be written as

$$\bar{\rho}|_{D_w} \cong \begin{bmatrix} \psi_1 & * \\ 0 & \psi_2 \end{bmatrix},$$

where $\psi_1 \oplus \psi_2|_{I_w}$ is isomorphic to the sum of the trivial character and $\bar{\chi}_3$.

- The image of the inertia group I_v in $\mathrm{GL}_2(\mathbb{F}_9)$ has odd order for every prime v of K over 5.

Then there exists a totally real number field F with F/K a solvable Galois extension, and a Hilbert-Blumenthal abelian variety A/F with real multiplication by $\mathcal{O} = \mathcal{O}_{\mathbb{Q}[\sqrt{5}]}$, such that

- The absolute ramification degree of F is odd at every prime of F over 3 and 5;
- A has good ordinary or multiplicative reduction at all primes of $F(\zeta_3)$ above 3 and 5;
- $A[\sqrt{5}]$ is an absolutely irreducible $\mathrm{Gal}(\bar{F}/F)$ -module;
- there exists a symplectic isomorphism of $\mathrm{Gal}(\bar{F}/F)$ -modules

$$\iota : A[3] \cong V_{\bar{\rho}}.$$

2 Proof of Proposition 1.3

Our main tool is an explicit description of the moduli space of HBAV's with real multiplication by $\mathcal{O}$ and full 3-level structure, worked out by Hirzebruch and van der Geer.

Lemma 2.1. *Let $S_{\mathbb{Z}}$ be the surface in $\mathbb{P}^5/\mathbb{Z}$ defined by the equations*

$$\sigma_1(s_0, \dots, s_5) = \sigma_2(s_0, \dots, s_5) = \sigma_4(s_0, \dots, s_5) = 0,$$

where σ_i is the i th symmetric polynomial. Note that $A_6 \cong \mathrm{PSL}_2(\mathbb{F}_9)$ acts on $S_{\mathbb{C}}$ by permutation of coordinates.

Let $X_{\mathbb{C}}$ be the Hilbert modular surface parametrizing complex HBAV's with real multiplication by $\mathcal{O}$ and full 3-level structure. Then $X_{\mathbb{C}}$ and $S_{\mathbb{C}}$ are related by a birational isomorphism which is compatible with the action of $\mathrm{PSL}_2(\mathbb{F}_9)$ on $X_{\mathbb{C}}$ and A_6 on $S_{\mathbb{C}}$.

Proof. [13, VIII.(2.6)] □

More precisely, the complement of the cusps in $X_{\mathbb{C}}$ maps to an open subvariety of $S_{\mathbb{C}}$. Write $Z_{\mathbb{C}} \subset S_{\mathbb{C}}$ for the complement of the image of this map. We write Q_1, Q_2, Q_4 for the hypersurfaces in $\mathbb{P}^5/\mathbb{Z}$ cut out by $\sigma_1 = 0, \sigma_2 = 0, \sigma_4 = 0$ respectively.

The morphism from $X_{\mathbb{C}}$ to $S_{\mathbb{C}}$ is defined as follows. Let Γ be the kernel of the reduction map $\mathrm{SL}_2(\mathcal{O}) \rightarrow \mathrm{SL}_2(\mathbb{F}_9)$. Then the complex vector space $S_2(\Gamma; \mathbb{C})$ of weight 2 cuspforms for Γ has dimension 5, and $\mathrm{PSL}_2(\mathbb{F}_9)$ acts on $S_2(\Gamma; \mathbb{C})$ through its irreducible 5-dimensional representation [13, VIII.2.5]. Let $s_0, \dots, s_5$ be a basis for $S_2(\Gamma; \mathbb{C})$ such that $\mathrm{PSL}_2(\mathbb{F}_9)$ acts by even permutations on $s_0, \dots, s_5$. Then taking $s_0, \dots, s_5$ as projective coordinates on $X_{\mathbb{C}}$ yields the rational morphism from $X_{\mathbb{C}}$ to $\mathbb{P}^5$ described above.

In particular, we have

$$\sigma_1(s_0, \dots, s_5) = \sigma_2(s_0, \dots, s_5) = \sigma_4(s_0, \dots, s_5) = 0.$$

Note that the other symmetric functions σ_3, σ_5 , and σ_6 in $s_0, \dots, s_5$ are cuspforms of weight 6, 10, 12 for the whole group $\mathrm{SL}_2(\mathcal{O})$. In section 2.2 we will describe these forms in terms of classical generators of the ring of level 1 modular forms.

The complex moduli space above descends to one defined over a ring of algebraic numbers. Let N be the product of all primes where $\bar{\rho}$ is ramified. (In particular, 3 divides N .) Then the representation $\bar{\rho}$ defines an étale group scheme $V_{\bar{\rho}}/\mathbb{Z}[1/N]$. The determinant condition on $\bar{\rho}$ yields a natural isomorphism $\wedge^2 V_{\bar{\rho}} \cong \mu_3 \otimes_{\mathbb{Z}} \mathcal{O}$.

Then there exists a proper scheme $\mathcal{X}^{\bar{\rho}}/\mathbb{Z}[1/N]$ parametrizing pairs (A, ι) , where A is a generalized HBAV, and

$$\iota : A[3] \cong V_{\bar{\rho}}$$

is an isomorphism of group schemes such that $\wedge^2 \iota$ is the identity map. Write $X^{\bar{\rho}}/K$ for the restriction of $\mathcal{X}^{\bar{\rho}}$ to $\mathrm{Spec} K$.

The surface $X^{\bar{\rho}}/K$ is geometrically isomorphic to X ; in particular, its geometric genus is still 5. Let $t_0, \dots, t_5$ be a set of sections spanning $H^0(X^{\bar{\rho}}, \Omega_{X^{\bar{\rho}}}^1)$. Then $t_0, \dots, t_5$ can be expressed as complex linear combinations of $s_0, \dots, s_5$. In particular, the rational map $X^{\bar{\rho}}/K \rightarrow \mathbb{P}^5/K$ defined by $t_0, \dots, t_5$ is a locally closed immersion of the complement of the cusps in $X^{\bar{\rho}}$ into $\mathbb{P}^5$ as an open subset in an intersection of smooth hypersurfaces of degrees 1, 2 and 4. Write $S^{\bar{\rho}}/K$ for the closure of the image of $X^{\bar{\rho}}$ in $\mathbb{P}^5$.

Note that there exists an isomorphism $\alpha : \mathbb{P}_{\mathbb{Q}}^5 \rightarrow \mathbb{P}_{\mathbb{Q}}^5$, depending on our choice of $t_0, \dots, t_5$, such that $S^{\bar{\rho}} = \alpha^{-1}S$. Write $Q_1^{\bar{\rho}}, Q_2^{\bar{\rho}}$ and $Q_4^{\bar{\rho}}$ for the hypersurfaces $\alpha^{-1}Q_1, \alpha^{-1}Q_2, \alpha^{-1}Q_4$. Then $S^{\bar{\rho}}$ is the complete intersection of $Q_1^{\bar{\rho}}, Q_2^{\bar{\rho}}$ and $Q_4^{\bar{\rho}}$, which are hypersurfaces in $\mathbb{P}^5/\mathbb{Q}$.

To prove Proposition 1.3, we will need to find a point on a twisted Hilbert modular variety $X^{\bar{\rho}}$ defined over a solvable extension of K . The geometric observation that enables us to find such points is the following. Let L/K be a line contained in the variety $Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}}$. Then $L \cap Q_4$ is a finite subscheme Σ of degree 4 of $S^{\bar{\rho}}$. Generically, Σ will split into 4 distinct points over a degree 4 (whence solvable!) extension of K . Now $Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}}$ is isomorphic to a quadric hypersurface in $\mathbb{P}^4$, so its Fano variety is rational. This means we have plenty of lines in $Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}}$, whence plenty of points in $S^{\bar{\rho}}$ defined over solvable extensions of K . What remains is to make sure we can find such points which satisfy the local conditions at 3, 5, and ∞ required in the theorem. Our strategy will be to define suitable lines over completions of K at the relevant primes, and finally to use strong approximation on the Fano variety $F_1(Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}})$ to find a global line which is adelically close to the specified local ones.

2.1 Archimedean places

Let c be a complex conjugation in $\mathrm{Gal}(\bar{K}/K)$, and let u be the corresponding real place of K .

The fact that $\bar{\rho}$ is odd implies that $\bar{\rho}(c)$ is conjugate to

$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

In particular, if

$$\bar{\rho}_0 = \begin{bmatrix} \bar{\chi}_3 & 0 \\ 0 & 1 \end{bmatrix},$$

we have

$$\bar{\rho}_0| \text{Gal}(\mathbb{C}/K_u) \cong \bar{\rho}| \text{Gal}(\mathbb{C}/K_u),$$

whence

$$S^{\bar{\rho}} \times_K K_u \cong S^{\bar{\rho}_0} \times_{\mathbb{Q}} K_u = S^{\bar{\rho}_0} \times_{\mathbb{Q}} \mathbb{R}.$$

If $s_0, \dots, s_5$ are our standard coordinates on S , we may take $t_0, \dots, t_5$ as coordinates on $S_{\mathbb{Q}}^{\bar{\rho}_0}$, where

$$(s_0, \dots, s_5) = (t_0 + \sqrt{-3}t_1, t_0 - \sqrt{-3}t_1, t_2 + \sqrt{-3}t_3, t_2 - \sqrt{-3}t_3, t_4, t_5). \quad (2.1.1)$$

Then $S_{\mathbb{Q}}^{\bar{\rho}_0}$ is isomorphic to the complete intersection

$$\begin{aligned} & \sigma_1(t_0 + \sqrt{-3}t_1, t_0 - \sqrt{-3}t_1, t_2 + \sqrt{-3}t_3, t_2 - \sqrt{-3}t_3, t_4, t_5) \\ &= \sigma_2(t_0 + \sqrt{-3}t_1, t_0 - \sqrt{-3}t_1, t_2 + \sqrt{-3}t_3, t_2 - \sqrt{-3}t_3, t_4, t_5) \\ &= \sigma_4(t_0 + \sqrt{-3}t_1, t_0 - \sqrt{-3}t_1, t_2 + \sqrt{-3}t_3, t_2 - \sqrt{-3}t_3, t_4, t_5) \\ &= 0. \end{aligned}$$

Now choose a real line $L_{\mathbb{R}}$ in $F_1(Q_1^{\bar{\rho}_0} \cap Q_2^{\bar{\rho}_0})(\mathbb{R})$ with the property that $L_{\mathbb{R}} \cap S^{\bar{\rho}_0}$ consists of four distinct real points. For instance, we may choose $L_{\mathbb{R}}$ to be the line

$$(t_0, t_1, t_2, t_3, t_4, t_5) = (7/15 + (4/3)t, -1, (4/5) - t, t, -2, -8/15 - (2/3)t).$$

Let L_u be the corresponding line in $F_1(Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}})(K_u)$.

2.2 Primes above 5

Let K_v be the completion of K at a prime v dividing 5, and let E_0 be the splitting field of $\bar{\rho}|G_{K_v}$. Note that, by hypothesis, $E_{v'}$ has odd absolute ramification degree.

As above, our aim is to find a suitable line in $(Q_1^{\bar{\rho}} \cap Q_2^{\bar{\rho}})$ over some unramified extension of $E_{v'}$. Since $\bar{\rho}$ is trivial on $\text{Gal}(\bar{\mathbb{Q}}_5/E_{v'})$, we may drop superscripts and look for lines on $Q_1 \cap Q_2$.

Lemma 2.2. *There exists a finite unramified extension E of $E_{v'}$ and a line L_v/E contained in $Q_1 \cap Q_2/E$ such that*

- L_v is disjoint from the cuspidal locus Z ;
- $(L_v \cap Q_4)(E)$ consists of 4 distinct E -points;
- For each $x \in (L_v \cap Q_4)(E)$, the functions

$$\sigma_5^{-6}(\sigma_3^2 - 4\sigma_6)^5$$

and

$$\sigma_5^{-3}\sigma_3^{-1}(\sigma_3^2 - 4\sigma_6)^3$$

have non-positive valuation when evaluated at x .

Proof. One checks that $Q_1 \cap Q_2$ is isomorphic over $\mathbb{Z}_5^{unr}$ to the Plücker quadric threefold $T := V(y_0y_1 + y_2y_3 + y_4^2) \subset \mathbb{P}^4$. We also know (see [4, §6, Ex. 22.6]) an explicit 3-parameter family of lines on T , which is to say a map

$$\lambda : \mathbb{P}^3 / \text{Spec } \mathbb{Z}_5 \rightarrow F_1(T);$$

moreover, λ is an isomorphism over any algebraically closed field. Composing λ with an isomorphism between T and $Q_1 \cap Q_2$ yields a map

$$L : \mathbb{P}^3 / \text{Spec } \mathbb{Z}_5^{unr} \rightarrow F_1(Q_1 \cap Q_2)$$

which is an isomorphism over any algebraically closed field.

The set of $\bar{\mathbf{p}} \in \mathbb{P}^3(\bar{\mathbb{F}}_5)$ such that $L(\bar{\mathbf{p}}) \cap Q_4/\bar{\mathbb{F}}_5$ consists of 4 distinct $\bar{\mathbb{F}}_5$ -points is Zariski-open. To check that it is not empty, we need only exhibit a single such line L in $(Q_1 \cap Q_2)/\bar{\mathbb{F}}_5$. One such line is

$$(s_0, s_1, s_2, s_3, s_4, s_5) = ((1 - \sqrt{-3})t, (1 + \sqrt{-3})t, -t + (1 + \sqrt{-3})u, -t + (1 - \sqrt{-3})u, t, -t - 2u).$$

One checks that the restriction of Q_4 to L is $-3t(8u^3 - t^3)$, which indeed has 4 distinct roots over $\bar{\mathbb{F}}_5$.

Let V be the closed subscheme of $S/\bar{\mathbb{F}}_5$ where the form $\sigma_3^2 - 4\sigma_6$ vanishes. Then V is a curve. Moreover, if x is a point in $S/\bar{\mathbb{F}}_5$, the subscheme of $\mathbb{P}^3/\bar{\mathbb{F}}_5$ parametrizing lines passing through x is one-dimensional. So the subscheme of $\mathbb{P}^3/\bar{\mathbb{F}}_5$ parametrizing lines intersecting V is at most two-dimensional. We may thus choose a point $\bar{\mathbf{p}} \in \mathbb{P}^3(\bar{\mathbb{F}}_5)$ such that $L(\bar{\mathbf{p}}) \cap Q_4/\bar{\mathbb{F}}_5$ consists of four distinct $\bar{\mathbb{F}}_5$ -points, none contained in V .

Now let $\mathbf{p}$ be a lift of $\bar{\mathbf{p}}$ to $\mathbb{P}^3(\mathbb{Q}_5^{nr})$. Then $L(\mathbf{p})$ is a line contained in $Q_1 \cap Q_2$ whose intersection with Q_4 consists of four distinct points defined over some unramified extension of $\mathbb{Q}_5$. Let E be the compositum of this extension with $E_{v'}$. Since Z is one-dimensional, we may choose $\mathbf{p}$ such that $L(\mathbf{p}) \cap Q_4$ is disjoint from Z , by the same argument as above.

Let x be a point in $L(\mathbf{p}) \cap Q_4$, and choose integral coordinates for x so that at least one coordinate has non-positive valuation. Then $(\sigma_3^2 - 4\sigma_6)(x)$ has non-positive valuation, so the third desired condition on $L(\mathbf{p})$ is satisfied. This completes the proof. $\square$

Now take L and E as in the lemma. Let x_1, x_2, x_3, x_4 be the four E -points making up $(L_v \cap S)(E)$. Then each x_i corresponds to an abelian variety A_i/E with real multiplication by $\mathcal{O}$ admitting an isomorphism $A[3] \cong \bar{\rho} \cong \mathbb{F}_9^{\oplus 2}$ of $\mathcal{O}$ -module schemes over E . Since $\mathcal{X}^{\bar{\rho}}/\mathcal{O}_E$ is a fine proper moduli space, we know that A_i extends to a semi-abelian scheme $\mathcal{A}_i/\mathcal{O}_E$.

We now want to show that each A_i has good ordinary or multiplicative reduction. We begin by recalling some facts on Hilbert modular forms for $\text{SL}_2(\mathcal{O})$. By a result of Nagaoka [8, Th. 2], the ring $M_{2*}(\text{SL}_2(\mathcal{O}), \mathbb{Z}[1/2])$ of even-weight symmetric modular forms over $\mathbb{Z}[1/2]$ for this group is generated by forms ϕ_2, χ_6 , and χ_{10} of weights 2, 6, 10. The form ϕ_2 is the weight 2 Eisenstein series, while χ_6 and χ_{10} are cuspforms.

By restricting to various modular curves on X , and comparing q -expansions, one can verify the following identities between the level 1 forms above and the symmetric functions σ_i in $s_0, \dots, s_5$:

$$\begin{aligned} \phi_2 &= -3\sigma_5^{-1}(\sigma_3^2 - 4\sigma_6) \\ \chi_6 &= \sigma_3 \\ \chi_{10} &= (-1/3)\sigma_5. \end{aligned}$$

(Note that the constants here depend on our original choice of the weight 2 forms s_i . Modifying that choice by a constant c would modify each formula above by $c^{k/2}$, where k is the weight of the modular form in the expression.)

So by our choice of L_v , the modular functions ϕ_2^3/χ_6 and ϕ_2^5/χ_{10} have non-positive valuation when evaluated on A_i . The desired ordinarity now follows from the following lemma.

Lemma 2.3. *Let A/E be an semi-HBAV over a finite extension $\mathcal{O}_E/\mathbb{Z}_5$. Suppose that the modular functions ϕ_2^3/χ_6 and ϕ_2^5/χ_{10} evaluated at A have non-positive valuation. Then A has potentially ordinary or multiplicative reduction.*

Proof. Let Ω be the determinant of the pushforward of the relative cotangent sheaf of $A/\mathcal{O}_E$. Then Ω is a free rank 1 $\mathcal{O}_E$ -module. Let ω be a section generating Ω . Then every modular form f with coefficients $\mathcal{O}_E$ has a well-defined value $f(A, \omega)$. Suppose $\phi_2(A, \omega) \in \mathfrak{m}_E$. Then by the hypothesis of the theorem, we have also that $\chi_6(A, \omega)$ and $\chi_{10}(A, \omega) \in \mathfrak{m}_E$. But this is impossible, as we show in the following paragraph.

Let $q \geq 7$ be a prime, let $\iota : (\mathbb{Z}/q\mathbb{Z})^2 \cong A[q]$ be an arbitrary full level q structure on A , and let f be a modular form of weight $2k$ and full level q . Since every even-weight modular form of full level q is integral over $M_{2*}(\Gamma(1), \mathbb{Z}) = \mathbb{Z}_5[\phi_2, \chi_6, \chi_{10}]$, we have $f(A, \iota, \omega) \in \mathfrak{m}_E$. But this is impossible, because for $2k$ sufficiently large, the sheaf $\Omega^{\otimes 2k}$ is very ample on the level q moduli scheme $\mathcal{X}(q)$.

We conclude that $\phi_2(A, \omega) \neq \mathfrak{m}_E$. So the mod 5 reduction $\bar{\phi}_2(\bar{A}, \bar{\omega})$ is not equal to 0. Since the reduction mod 5 of ϕ_2^2 is the Hasse invariant, A has good ordinary or multiplicative reduction by [1]. $\square$

In order to keep straight the primes of K over 5, we now give the extension $E/E_{v'}$ the new name $E'_{v'}/E_{v'}$.

2.3 Primes above 3

We begin by observing that we can apply to $\bar{\rho}$ a global quadratic twist ψ such that

$$\bar{\rho}|D_w \cong \begin{bmatrix} \bar{\chi}_3 & * \\ 0 & 1 \end{bmatrix} \quad (2.3.2)$$

for all primes w of K dividing 3. Since twisting does not affect modularity, we may assume that (2.3.2) holds for our original $\bar{\rho}$.

Let w be a prime of K dividing 3, and let K_w be the completion of K and w . Now the $*$ in (2.3.2) is a cocycle corresponding to an element $\lambda \in K_w^* \otimes_{\mathbb{Z}} \mathbb{F}_9$. Write $\bar{\rho}_\lambda$ for the representation $\bar{\rho}|G_w$.

Lemma 2.4. *There exists a line L_w in $\mathbb{P}_{K_w}^5$ satisfying the following conditions.*

- L_w is contained in $Q_1^{\bar{\rho}_\lambda} \cap Q_2^{\bar{\rho}_\lambda}$.
- The intersection $L_w \cap Q_4^{\bar{\rho}_\lambda}$ splits into four distinct points over an unramified extension $E'_{w'}$ of K_w .
- The four HBAV's A_1, A_2, A_3, A_4 corresponding to the four points of $L_w \cap Q_4^{\bar{\rho}_\lambda}(\mathbb{Q}_3^{nr})$ have potentially multiplicative reduction.

Proof. We define new coordinates $x_0, \dots, x_5$ on $\mathbb{P}^5$ by the rule

$$\begin{aligned} x_0 &= \omega s_0 + \omega^2 s_1 + s_4 \\ x_1 &= \omega^2 s_0 + \omega s_1 + s_4 \\ x_4 &= s_0 + s_1 + s_4 \\ x_2 &= \omega s_2 + \omega^2 s_3 + s_5 \\ x_3 &= \omega^2 s_2 + \omega s_3 + s_5 \\ x_5 &= s_2 + s_3 + s_5 \end{aligned}$$

where ω is a cube root of unity. With these coordinates, one checks that $Q_1^{\bar{\rho}^\lambda}$ is defined by $x_4 + x_5$ and $Q_2^{\bar{\rho}^\lambda}$ by

$$x_4^2 + x_5^2 - x_0 x_1 - x_2 x_3 + 3x_4 x_5.$$

So a family of lines in $Q_1^{\bar{\rho}^\lambda} \cap Q_2^{\bar{\rho}^\lambda}$ is given by

$$L_{a,b,c} : x_0 = ax_2 + bx_4, x_3 = -ax_1 + cx_4, x_4 = -(bx_1 + cx_2), x_5 = -x_4.$$

One checks that the equation for $Q_4^{\bar{\rho}^\lambda}$ is given by

$$\begin{aligned} -3x_0 x_1 x_4 x_5 - 3x_2 x_3 x_4 x_5 + 3x_0 x_1 x_2 x_3 + x_4 x_5 (x_4^2 + 3x_4 x_5 + x_5^2) - 3x_0 x_1 x_5^2 - 3x_2 x_3 x_4^2 \\ + \lambda_1^3 x_0^3 x_5 + \lambda_1^{-3} x_1^3 x_5 + \lambda_2^3 x_2^3 x_4 + \lambda_2^{-3} x_3^3 x_4 \end{aligned}$$

where λ_1, λ_2 are elements of K whose class in $K^*/(K^*)^3$ is determined by λ .

The equation for $Q_4^{\bar{\rho}^\lambda}$ restricted to $L_{a,b,c}$ is of the form

$$P = \sum_{i=0}^4 P_i(a, b, c) x_1^i x_2^{4-i}.$$

Suppose that $\text{ord}_w(b)$ and $\text{ord}_w(c)$ are approximately equal and that both are much greater than $\text{ord}_w(a)$, which is in turn much greater than 0. Then one checks that

$$\begin{aligned} P_4(a, b, c) &= \lambda_1^{-3} b + \text{higher order terms} \\ P_3(a, b, c) &= \lambda_1^{-3} c + \text{higher order terms} \\ P_2(a, b, c) &= -3a^2 + \text{higher order terms} \\ P_1(a, b, c) &= -\lambda_2^3 b + \text{higher order terms} \\ P_0(a, b, c) &= -\lambda_2^3 c + \text{higher order terms.} \end{aligned}$$

It is an easy calculation that P then factors over K_w^{unr} into a constant and two quadratics, one of which has discriminant with valuation equal to that of $\lambda_1^{-3} a^{-2} b$, the other of which has discriminant with valuation equal to that of $\lambda_2^3 a^{-2} c$. In particular, if b and c are chosen so that $\text{ord}_w(b)$ has the same parity as $\text{ord}_w(\lambda_1)$, and $\text{ord}_w(c)$ the same parity as $\text{ord}_w(\lambda_2)$, the points of $L_{a,b,c} \cap Q_4^{\bar{\rho}^\lambda}$ are defined over an unramified extension of K_w .

Note that since $\text{ord}_w(a), \text{ord}_w(b)$, and $\text{ord}_w(c)$ are much larger than 0, we know that $L_{a,b,c}$ is very v -adically close to the cusp line $L_{0,0,0}$. Moreover, the form for P given above shows that the four points in $L_{a,b,c} \cap Q_4^{\bar{\rho}^\lambda}$ are w -adically close to $[0 : 1 : 0 : 0 : 0 : 0]$ and $[0 : 0 : 1 : 0 : 0 : 0]$.

These points do not lie in the image of any non-cuspidal point of the Hilbert modular surface X [13, VIII(2.6)]. So the abelian varieties A_1, A_2, A_3, A_4 corresponding to these four points to the points of $L_{a,b,c} \cap Q_4^{\bar{\rho}_\lambda}$ have potentially multiplicative reduction at w , as desired. $\square$

The fact that A_i has potentially multiplicative reduction implies that the ℓ -adic representation $\rho_{A_i, \ell}$ attached to A_i satisfies

$$\rho_{A_i, \ell}|_{I_{w'}} \cong \begin{bmatrix} * & * \\ 0 & * \end{bmatrix}.$$

The diagonal entries are finite-order characters with image in $\mathcal{O}^*$, which must have image in ± 1 . It follows that either A_i or its quadratic twist $A_i \otimes \bar{\chi}_3$ has semi-stable reduction at w' . Note that $A_i[3](\bar{\mathbb{Q}}_3)$ contains a canonical subgroup A_i^0 , consisting of those points reducing to the identity in the Néron model of A_i . Now the action of $I_{w'}$ on $A_i[3]$ is isomorphic to

$$\begin{bmatrix} \bar{\chi}_3 & * \\ 0 & 1 \end{bmatrix},$$

and $A_i[3]$ has semistable reduction if and only if the canonical subgroup is the subgroup on which $I_{w'}$ acts as $\bar{\chi}_3$. Note that $A_i[3]$ is semistable whenever $*$ is nonzero.

Each of the points in $L_{a,b,c} \cap Q_4^{\bar{\rho}_\lambda}$ corresponds to an abelian variety A_i together with an isomorphism ϕ_i from $A_i[3]$ to $V_{\bar{\rho}_\lambda}$. The (A_i, ϕ_i) correspond to points of $X(\bar{\mathbb{Q}}_3)$ lying in a small 3-adic neighborhood of a cusp; since the canonical subgroup varies 3-adically continuously, the image $\phi_i(A_i^0)$ is the same subgroup of $V_{\bar{\rho}_\lambda}$ for each i . In particular, either all four of the A_i have semistable reduction at w' , or all four have semistable reduction after twisting by $\bar{\chi}_3$.

2.4 The global construction

We now combine the local arguments above into the global statement we desire.

Choose a finite Galois extension K'/K such that;

- K' is totally real;
- K'/K is solvable;
- The completion of K' at any prime v above 5 is isomorphic to an unramified extension of E'_v ;
- The completion of K' at any prime w above 3 is isomorphic to an unramified extension of E'_w ;
- $F_1((Q_1 \cap Q_2)^{\bar{\rho}})$ is rational over K' . (Since $F_1((Q_1 \cap Q_2)^{\bar{\rho}})$ is geometrically rational, this amounts to trivializing an element of the Brauer group.)

(See [12, Lemma 2.2] for the existence of K' .)

Since $F_1((Q_1 \cap Q_2)^{\bar{\rho}})$ is a rational variety, we can choose $L \in F_1((Q_1 \cap Q_2)^{\bar{\rho}})(K')$ such that the image of L under the map

$$F_1((Q_1 \cap Q_2)^{\bar{\rho}})(K') \rightarrow \bigoplus_{v_i|5} F_1((Q_1 \cap Q_2)^{\bar{\rho}})(K'_{v_i}) \oplus \bigoplus_{w_i|3} F_1((Q_1 \cap Q_2)^{\bar{\rho}})(K'_{w_i}) \oplus \bigoplus_{u|\infty} \mathbb{F}_1((Q_1 \cap Q_2)^{\bar{\rho}})(K'_u)$$

is arbitrarily adelicly close to $(L_{v_1}, \dots, L_{w_1}, \dots, L_{u_1}, \dots)$.

The intersection $L \cap S^{\bar{\rho}}$ is a zero-dimensional scheme of degree 4 over K' . Modifying our choice of L if necessary, we can arrange for $L \cap S$ to be in the image of the rational map from $X^{\bar{\rho}}$. Let F be a splitting field for $L \cap S^{\bar{\rho}}$. Note that F is solvable over K' , whence also over K . Then we can think of $L \cap S$ as specifying four HBAV's A_i/F , with $A_i[3]/F \cong V_{\bar{\rho}}/F$.

By our choices of L_u , the field F is totally real. Similarly, our choices of L_v and L_w guarantee that A_i and F satisfy the local conditions at 3 and 5 stated in the theorem, by a theorem of Kisin [7].

It remains only to check that L can be chosen so that $A_i[\sqrt{5}]$ is an absolutely irreducible $\text{Gal}(\bar{F}/F)$ -module, for some i . Let $\pi : T_{\mathbb{C}} \rightarrow F_1(Q_1 \cap Q_2)_{\mathbb{C}}$ be the degree 4 cover defined by $\pi^{-1}(L) = (L \cap S)(\mathbb{C})$. Then T projects to S , and the covering of S by the Hilbert modular surface for the congruence subgroup of full level $\sqrt{5}$ pulls back to a covering $T' \rightarrow T$. The covering T'/T is Galois with group $\text{PSL}_2(\mathbb{F}_5)$, and the Galois group of the cover $T'/F_1(Q_1 \cap Q_2)$ is thus a subgroup G of the wreath product of S_4 and $\text{PSL}_2(\mathbb{F}_5)$.

We want to show that G is large. Let B be an HBAV over a number field M such that the map

$$\text{Gal}(\bar{\mathbb{Q}}/M) \rightarrow \text{GL}(B[3]) \oplus \text{GL}(B[\sqrt{5}])$$

is surjective, and let p be a point of $S(M(B[3]))$ corresponding to B . Let ℓ be a line in $Q_1 \cap Q_2$ passing through p ; we can choose ℓ to be defined over a quadratic extension of M . Now the fact that the map

$$\text{Gal}(\bar{\mathbb{Q}}/M(B[3])) \rightarrow \text{GL}(B[\sqrt{5}])$$

is surjective implies that the Galois group G contains a copy of $\text{PSL}_2(\mathbb{F}_5)$. By Ekedahl's version of Hilbert's irreducibility theorem [3], we can now choose L in the adelic neighborhood specified above in such a way that one of the four abelian varieties A_i/F has $A_i[\sqrt{5}]$ an absolutely irreducible $\text{Gal}(\bar{F}/F)$ -module. This completes the proof of Proposition 1.3.

3 Modularity

Now that we have exhibited $\bar{\rho}$ as a representation appearing on the torsion points of an abelian variety, we can prove that $\bar{\rho}$ is modular. Our argument proceeds exactly along the lines of [12] and [6]. We begin by recording the case we need of a theorem of Skinner and Wiles.

Theorem 3.1. *Let K be a totally real number field, let $p > 2$ be a rational prime, let L be a finite extension of $\mathbb{Q}_p$, and let*

$$\rho : \text{Gal}(\bar{K}/K) \rightarrow \text{GL}_2(L)$$

be a continuous irreducible representation ramified at only finitely many primes. Suppose

- $\det \rho = \psi \chi_p^{k-1}$ for some finite-order character ψ ;
- ρ is ordinary in the sense that, for each prime v of K dividing p ,

$$\rho|_{D_v} \cong \begin{bmatrix} \chi_p & * \\ 0 & 1 \end{bmatrix};$$

- ρ is D_v -distinguished for all primes v of K dividing p ;
- There exists an ordinary modular Galois representation ρ' and an isomorphism between the mod p representations $\bar{\rho}$ and $\bar{\rho}'$.

Then ρ is modular.

Proof. See [11]. □

We are now ready to prove our main result.

Theorem 3.2. *Let*

$$\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{F}_9)$$

be an odd, absolutely irreducible Galois representation such that

- *The restriction of $\bar{\rho}$ to D_3 can be written as*

$$\bar{\rho}|_{D_3} \cong \begin{bmatrix} \psi_1 & * \\ 0 & \psi_2 \end{bmatrix},$$

where ψ_1 and ψ_2 are distinct characters such that $(\psi_1 \oplus \psi_2)|_{I_3}$ is isomorphic to the direct sum of the trivial character and $\bar{\chi}_3$.

- *The image of the inertia group I_5 lies in $\text{SL}_2(\mathbb{F}_9)$, and has odd order.*

Then $\bar{\rho}$ is modular.

Proof. First of all, $\theta = (\det \bar{\rho})^{-1} \bar{\chi}_3$ is a character of Galois which annihilates complex conjugation, since $\bar{\rho}$ is odd. We thus have a totally real abelian extension $K/\mathbb{Q}$ defined by $\text{Gal}(\bar{K}/K) = \ker \theta$. Since $\det(\bar{\rho})(I_5)$ is trivial, I_5 lies in the kernel of θ , and K is unramified at 5. Likewise, $\theta(I_3)$ is trivial, so K is unramified at 3.

By applying a quadratic twist, we may assume that

$$\bar{\rho}|_{D_3} \cong \begin{bmatrix} \bar{\chi}_3 & * \\ 0 & 1 \end{bmatrix}.$$

Now the conditions on $\bar{\rho}|_{D_3}$ and $\bar{\rho}|_{D_5}$ imply the corresponding local conditions in Proposition 1.3. We may now choose an extension F_0/K and an abelian variety A/F_0 satisfying the four hypotheses given in that Proposition.

From here, we proceed along the lines of [12]. First, we claim that the irreducible representation

$$\bar{\rho}_{A, \sqrt{5}} : \text{Gal}(\bar{\mathbb{Q}}/F_0) \rightarrow \text{GL}_2(\mathbb{F}_5)$$

induced by the torsion subscheme $A[\sqrt{5}]$ is modular.

Now it follows from the discussion in section 2.3 that either A or $A \otimes \bar{\chi}_3$ has multiplicative reduction at all primes of F_0 over 3. In case it is $A \otimes \bar{\chi}_3$ which is semistable, return to the beginning, replace $\bar{\rho}$ by $\bar{\rho} \otimes \bar{\chi}_3$, and start over. We may now assume A has multiplicative reduction at all primes of F_0 over 3.

Now the subgroup

$$\bar{\rho}_{A, \sqrt{5}}(I_w) \subset \text{SL}_2(\mathbb{F}_5)$$

is unipotent. Thus, we can find a totally real solvable extension F/F_0 , unramified over 5 and with odd ramification degree at every prime over 3, such that D_w acts trivially on $A[\sqrt{5}]$ for every prime w of F dividing 3. Then the twist of the modular curve $X(5)_F$ by $\bar{\rho}_{A, \sqrt{5}}$ is isomorphic to $X(5)$ when

base changed to any 3-adic completion of F . In particular, there exists an elliptic curve E/F such that $E[5]/F \cong A[\sqrt{5}]/F$ and E has good ordinary reduction at each prime of F over 3. By another use of Ekedahl [3], we can assume that $E[3]$ is an absolutely irreducible $\text{Gal}(\bar{\mathbb{Q}}/F)$ -module.

Since F has odd absolute ramification degree at all primes over 3, we have by Proposition 1.2 that $E[3]/F$ is distinguished at all primes over 3. The mod-3 representation $E[3]$ is modular by the Langlands-Tunnell theorem, and the 3-adic Galois representation T_3E is modular by Theorem 3.1. It follows that T_5E is modular, and so $E[5]$, whence also $A[\sqrt{5}]/F$, is modular.

By hypothesis, A has good ordinary or multiplicative reduction at 5, so $T_{\sqrt{5}}A$ is an ordinary representation. Because $F/\mathbb{Q}$ has odd ramification degree over 5, $T_{\sqrt{5}}A$ is D_v -distinguished for all primes v dividing 5. Now $T_{\sqrt{5}}A/F$ is modular by another application of Theorem 3.1. This implies that T_3A/F is also modular, which in turn implies the modularity of $A[3]/F$, which is the restriction to $\text{Gal}(\bar{F}/F)$ of our original representation $\bar{\rho}$.

Recall that the restriction of $\bar{\rho}$ to the decomposition group D_3 is of the form

$$\bar{\rho}|_{D_3} \cong \begin{bmatrix} \bar{\chi}_3 & * \\ 0 & 1 \end{bmatrix}.$$

By results of Ramakrishna as refined by Taylor [12, Thm 1.3], $\bar{\rho}$ can be lifted to a 3-adic representation

$$\rho : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(W(\mathbb{F}_9))$$

such that

$$\rho|_{D_3} \cong \begin{bmatrix} \chi_3 & * \\ 0 & 1 \end{bmatrix}.$$

Now $\rho|_{\text{Gal}(\bar{F}/F)}$ is an ordinary, D_3 -distinguished 3-adic representation of $\text{Gal}(\bar{F}/F)$ whose reduction mod 3 is isomorphic to the modular representation $A[3]$. Applying Theorem 3.1 once more, using T_3A as ρ' , we have that $\rho|_{\text{Gal}(\bar{F}/F)}$ is modular. Now we argue by cyclic descent as in [12]. Let F' be a subfield of F such that F/F' is a cyclic Galois extension. Then the automorphic form π on $\text{GL}_2(F)$ corresponding to ρ is preserved by $\text{Gal}(F/F')$. Therefore, π descends to an automorphic form on $\text{GL}_2(F')$. Continuing inductively, one finds that ρ itself is associated to a modular form on $\text{GL}_2(\mathbb{Q})$; therefore, its mod 3 reduction $\bar{\rho}$ is modular. $\square$

This case of Serre's conjecture can be used to prove the modularity of Hilbert-Blumenthal abelian surfaces, under some conditions on reduction at 3 and 5.

Corollary 3.3. *Let $A/\mathbb{Q}$ be a Hilbert-Blumenthal abelian surface which has good ordinary reduction at 3 and 5. Then A is a quotient of $J_0(N)$ for some integer N .*

Proof. Let v be a prime of the field of real multiplication dividing 3. If $A[v]$ is absolutely reducible, then the corollary follows from a theorem of Skinner and Wiles [10]. So we may assume that $A[v]$ is absolutely irreducible.

If 3 is split or ramified in the ring of real multiplication $\mathcal{O}$, then the corollary follows from Langlands-Tunnell applied to $A[v]$ followed by Diamond's refinement of the theorem of Wiles and Taylor-Wiles [2]. If, on the other hand, 3 is inert in A , then $A[3]$ yields a representation $\rho_{A,3} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{F}_9)$, which is easily seen to satisfy the conditions of Theorem 3.2. So $A[3]$ is modular, and it follows, again by Diamond's theorem, that A is modular. $\square$

References

- [1] F. Andreatta and E.Z. Goren. Hilbert modular forms: mod p and p -adic aspects. Preprint.
- [2] F. Diamond. On deformation rings and Hecke rings. *Ann. Math.* 144 (1996), 137–166.
- [3] T. Ekedahl. An effective version of Hilbert’s irreducibility theorem. In *Séminaire de Théorie des Nombres, Paris 1988-1989*, Progr. Math. 91, pages 241–249. Birkhäuser Boston, Boston, 1990.
- [4] J. Harris. *Algebraic Geometry*. Springer-Verlag, 1992.
- [5] F. Hirzebruch. The ring of Hilbert modular forms for real quadratic fields of small discriminant. In *Modular Functions of One Variable VI*, number 627 in Lecture Notes in Mathematics, pages 288–323. Springer-Verlag, 1977.
- [6] C. Khare. Mod p descent for Hilbert modular forms. *Math. Res. Lett.* 7 (2000), no. 4, 455–462.
- [7] M. Kisin. Local constancy in p -adic families of Galois representations. *Math. Z.* 230 (1999), no. 3, 569–593.
- [8] S. Nagaoka. On the ring of Hilbert modular forms over $\mathbf{Z}$. *J. Math. Soc. Japan* 35 (1983), 547–555.
- [9] N. Shepherd-Barron and R. Taylor. Mod 2 and mod 5 icosahedral representations. *J. Amer. Math. Soc.* 10(2):283–298, 1997.
- [10] C. Skinner and A. Wiles. Residually reducible representations and modular forms. *Inst. Hautes Études Sci. Publ. Math.* 89 (1999), 5–126 (2000)
- [11] C. Skinner and A. Wiles. Nearly ordinary deformations of irreducible residual representations. To appear, *Ann. Fac. Sci. Toulouse Math.* (6)
- [12] R. Taylor. On icosahedral Artin representations II. Preprint. Available at <http://www.math.harvard.edu/~rtaylor>.
- [13] G. van der Geer. *Hilbert modular surfaces*. Springer-Verlag, 1988.